



Czech ACM Student Chapter
University of West Bohemia in Pilsen
Masaryk University
Pavol Jozef Šafárik University in Košice

Czech Technical University in Prague
Technical University of Ostrava
University of Žilina
Comenius University



CTU Open Contest 2026

Accident

`accident.(c|cpp|py), Accident.(java|kt)`

Míša was walking home from school when he tripped on the pavement and dropped his right-angled triangle on the ground. The fall damaged the triangle, and it may no longer be right-angled. Now he wants to fix it with minimal effort — by moving just one of its vertices so that it becomes right-angled again.

Help Míša decide the minimum distance he needs to move one of the vertices, so the triangle becomes right-angled again. It is guaranteed that the input points are not collinear (i.e., they do not lie on a single line) and no two points are the same.

Input Specification

The input consists of three lines, each describing one vertex of the triangle. Each line contains two space-separated integers x and y ($-100 \leq x, y \leq 100$), the coordinates of a point.

Output Specification

Output a decimal number, the minimum distance that one of the points needs to be moved so that this point and the other two points form the vertices of a right-angled triangle.

Your answer will be considered correct if its absolute error does not exceed 10^{-5} .

Sample Input 1

4 0
0 4
-1 -1

Output for Sample Input 1

1.414213562


Sample Input 2

10 13
14 -4
-42 31

Output for Sample Input 2

9.013726194

In the first sample the cheapest fix is to slide the point $(-1, -1)$ by $\sqrt{2} \approx 1.41421$ towards the origin, where it forms a right angle with $(4, 0)$ and $(0, 4)$.



Sample Input 1